

Historic, archived document

Do not assume content reflects current scientific knowledge, policies, or practices.

The Forest Biomass Resource of the United States



United States
Department of
Agriculture

Forest Service

General
Technical
Report WO-57

September 1990

The Forest Biomass Resource of the United States

Noel D. Cost, Project Leader, SE¹
James O. Howard, Project Leader, PNW
Bert Mead, Research Forester, PNW
William H. McWilliams, Research Forester, SO
W. Brad Smith, Research Forester, NC
Dwane D. Van Hooser, Project Leader, INT
Eric H. Wharton, Research Forester, NE

¹Forest Service experiment stations are: SE=Southeastern; PNW=Pacific Northwest; SO=Southern; NC=North Central; INT=Intermountain; NE=Northeastern.

Acknowledgements

The authors would like to thank Daniel D. Oswald and Karen Waddell--Pacific Northwest Forest and Range Experiment Station, Portland, OR--for their assistance with the Resources Planning Act data base and the information retrieval system that supports it; and James T. Bones--Forest Inventory, Economics, and Recreation Research, Washington, DC--for his contributions to the development of this study.

Highlights

- The dry weight of all live trees on timberland is almost 18 billion dry tons, or 36.9 dry tons per acre.
- Tree biomass is evenly divided between the two major species groups -- softwoods accounting for 48 percent and hardwoods for 52 percent.
- Most of the softwood biomass--more than 5 billion dry tons, or 61 percent of the softwood tree biomass--is in the West. Approximately 9 billion dry tons, or 93 percent of the hardwood tree biomass, is in the East.
- The largest portion of total tree biomass, almost 72 percent, is found in tree boles. Tree tops account for 17 percent, and cull trees 11 percent.
- Most of the total tree biomass is on privately owned land. Ten billion dry tons, or 55 percent of the total tree biomass, is on lands owned by individuals and corporations (other than forest industry).
- Almost 3 billion dry tons of total tree biomass, or 81 percent of all national forest biomass, is on national forests in the West. Two billion dry tons of biomass on forest industry lands, or 77 percent of all forest industry biomass, is concentrated in the South and West.
- Information on additional components of total forest biomass varies by region. These components can contribute significant amounts of biomass. Hardwood tree biomass in the Lake States, for example, is 73 percent of the total hardwood forest biomass. The remainder is stumps, foliage, seedlings, saplings, and shrubs.

Background

Over the last decade, biomass statistics have been published for most states. However, the existing aggregate data are either limited or out of date. The most recent statistics on biomass were for 1980 (U.S. Department of Agriculture 1981). The development of such data continues to lag even though user interest is high. This study was initiated to provide current biomass data for the United States and was conducted by Forest Service research units throughout the country.

Initially, biomass statistics were used primarily to determine the amount of logging residue that might be generated from commercial timber harvesting. Biomass statistics were reported according to timber-oriented components of the total forest resource: wood and bark weight, biomass of the tree bole, biomass on timberland, etc. Many of the conventional definitions of tree biomass are still slanted toward industrial wood use.

However, recent policy is being made at the State, national, and international level on the optimum allocation of the total forest resource for a variety of traditional and nontraditional uses. New segments of the research community have begun to study the global effects of climate change and the effect of manmade pollution on major ecosystems of the world. Current biomass standards may have little application for environmental assessments and global resource analyses.

The biomass statistics presented in this report remain focused on standards related to the timber component of the forest resource, since most users of forest resource data continue to think in terms of timber production. For the first time, though, additional data has been compiled on the total forest resource, which includes all forest land areas regardless of productivity potential. It also includes stumps, foliage, seedlings, saplings, and shrubs.

The biomass statistics we developed were based on current data. Under the authority of the Forest and Rangeland Renewable Resources Planning Act of 1974 and the Forest

and Rangeland Renewable Resources Research Act of 1978, the USDA Forest Service conducts periodic forest inventories of all States. We used the most current biomass data from these forest inventories.

The data was compiled from existing data bases. Where no data existed, volume data compiled from the timber chapter of the 1989 Resource Planning Act assessment were used as a base, applying appropriate conversion factors to estimate bole biomass. Total tree biomass estimates were developed through two methods: one from integrated biomass equations and the other from component estimates that were added to the information from the data base. Additional components for both methods were provided as available.

Definition of Terms

Biomass

Forest biomass. The weight of wood, bark, and foliage in all living woody plants above the ground, measured in oven-dry tons. Includes tree biomass; stump and foliage biomass associated with tree biomass; and the biomass of seedlings, saplings, and shrubs. Does not include the biomass of grasses, forbs, etc.

Tree biomass. The weight of wood and bark in all living trees 5.0 inches in diameter at breast height and larger above a 1-foot stump height, measured in oven-dry tons. Includes the biomass of live sound trees and the biomass of live cull trees.

Forest Components

Commercial tree species. Softwood and hardwood tree species suitable for industrial wood products. Excludes tree species of typically small size, poor form, or inferior quality.

Foliage. The leaves and fruits of live sound and live cull trees, 5.0 inches in diameter at breast height and larger.

Hardwoods. Dicotyledonous trees, usually broad-leaved and deciduous.

Live sound trees. Live trees of commercial species 5.0 inches in diameter at breast height and larger, above a 1-foot stump height, and meeting regional specifications for freedom from defect. Includes tree boles and tree tops.

Live cull trees. Live trees 5.0 inches in diameter at breast height and larger, above a 1-foot stump height, not meeting regional specifications for freedom from defect. This includes live trees of commercial tree species that have such poor form that they do not contain a sawlog, that are not commercial tree species, and in which 50 percent of the cull is rotten.

Saplings. Live trees between 1.0 inch and 4.9 inches in diameter at breast height.

Seedlings. Live trees less than 1.0 inch in diameter at breast height that are free to grow.

Shrubs. Plants that have persistent, woody stems and relatively low growth habits, and that generally produce several basal shoots instead of a single bole. They differ from trees by their low stature and nonarborescent form (Society of Range Management 1974).

Softwoods. Coniferous trees, usually evergreen and having needles or scalelike leaves.

Stumps. The central stem of live sound and live cull trees, 5.0 inches in diameter at breast height and larger, between the ground and a 1-foot height.

Tree boles. The central stem of live trees, 5.0 inches in diameter at breast height and larger, between a 1-foot stump height and a 4-inch top diameter outside the bark, or until the stem breaks into branches before it reaches this minimum diameter.

Tree tops. The branches of live sound and live cull trees, 5.0 inches in diameter at breast height and larger, outside of the minimum top diameter for tree boles. Includes the upper stem, limbs, twigs, etc.

Land Classes

Forest industry lands. Lands owned by companies or individuals that operate primary wood-using facilities.

Forest land. Land areas 1 acre in size at least 10-percent stocked with trees of any size. Does not include productive-reserved land.

National forest lands. Federal lands legally designated as national forests or purchase units.

Other forest land. Forest land not producing or capable of producing at least 20 cubic feet of annual crops of commercial timber. Does not include productive-reserved land.

Other private lands. All privately owned lands except forest industry lands.

Other public lands. Lands other than national forests that are administered by Federal, State, county, or local municipalities.

Productive-Reserved land. Forest land withdrawn from timber utilization because of administrative designation, such as park land.

Timberland. Forest land producing or capable of producing at least 20 cubic feet of annual crops of commercial timber. Does not include productive-reserved land.

Literature Cited

Society for Range Management, Range Term Committee, M.M. Kothmann, Chair. 1974. A Glossary of Terms Used in Range Management. Denver, CO: Soc. for Range Management: 25.

U.S. Department of Agriculture. 1981. Tree Biomass--a state-of-the art compilation. Gen. Tech. Rep. WO-33; Washington, DC: U.S. Department of Agriculture, Forest Service. 34 p.

Index to Tables

The data are presented in tables of tree biomass by class of timber and ownership; forest biomass by class of material—including stumps, foliage, seedlings, saplings, and shrubs; and tree biomass per acre on forest land. Because most of the data were developed from the timber assessment volume data in the 1989 Resources Planning Act, the only species information is major species group (softwoods and hardwoods).

The statistics are reported in dry weight. Biomass is often reported in green weight, a unit desirable to the timber-using industry since it reflects the economic cost of removing wood from the forest. Dry weight is a more common base. It yields an easier solution to environmental questions, such as the carbon content of the world's standing timber, and is more consistent over time, avoiding seasonal variations that occur when measuring in green weight.

| Table | | Page |
|--------------|---|-------------|
| 1 | Softwood tree biomass on timberland in the United States, by State and section, and class of timber, 1987 | 4 |
| 2 | Hardwood tree biomass on timberland in the United States, by State and section, and class of timber, 1987 | 6 |
| 3 | Total tree biomass on timberland in the United States, by State and section, and class of timber, 1987 | 8 |
| 4 | Softwood tree biomass on timberland in the United States, by State and section, and ownership class, 1987 | 10 |

| Table | | Page |
|--------------|--|-------------|
| 5 | Hardwood tree biomass on timberland in the United States, by State and section, and ownership class, 1987 | 12 |
| 6 | Total tree biomass on timberland in the United States, by State and section, and ownership class, 1987 | 14 |
| 7 | Total forest biomass on timberland in the United States, by major species group, section, region, and class of material, 1987 | 16 |
| 8 | Total tree biomass per acre on forest land in the United States, by major species group, section, region, and land class, 1987 | 19 |

Note: Values in the tables may not add due to rounding.

Table 1.--Softwood tree biomass on timberland in the United States, by State and section,
and class of timber, 1987

| State and section | Live sound trees | | | Cull trees | Total tree biomass |
|-------------------------|--------------------------------|-----------|---------|---------------|-----------------------|
| | Tree boles | Tree tops | Total | | |
| | ----- (Million dry tons) ----- | | | | |
| Connecticut | 7.2 | 2.4 | 9.6 | 0.5 | 10.1 |
| Maine | 219.8 | 85.3 | 305.1 | 30.6 | 335.7 |
| Massachusetts | 28.5 | 9.3 | 37.9 | 2.2 | 40.1 |
| New Hampshire | 56.3 | 20.2 | 76.5 | 6.8 | 83.3 |
| Rhode Island | 0.8 | 0.3 | 1.1 | (a) | 1.1 |
| Vermont | 31.3 | 11.5 | 42.9 | 7.8 | 50.7 |
| Total, New England | 344.0 | 129.0 | 473.0 | 47.9 | 520.9 |
| Delaware | 2.5 | 0.9 | 3.4 | (a) | 3.4 |
| Maryland | 12.3 | 4.6 | 16.8 | 0.1 | 17.0 |
| New Jersey | 9.6 | 3.8 | 13.3 | 2.0 | 15.3 |
| New York | 101.5 | 33.2 | 134.7 | 10.1 | 144.8 |
| Pennsylvania | 32.2 | 10.0 | 42.2 | 0.4 | 42.6 |
| West Virginia | 16.2 | 5.8 | 22.0 | 1.8 | 23.8 |
| Total, Middle Atlantic | 174.3 | 58.2 | 232.5 | 14.5 | 246.9 |
| Michigan | 84.8 | 18.2 | 103.0 | 4.8 | 107.8 |
| Minnesota | 53.9 | 12.1 | 66.0 | 4.5 | 70.6 |
| North Dakota | (a) | (a) | (a) | (a) | (a) |
| South Dakota | 23.6 | 3.1 | 26.6 | 0.3 | 26.9 |
| Wisconsin | 55.1 | 10.4 | 65.5 | 2.7 | 68.1 |
| Total, Lake States | 217.4 | 43.8 | 261.1 | 12.3 | 273.4 |
| Illinois | 1.8 | 0.3 | 2.1 | 0.1 | 2.2 |
| Indiana | 3.0 | 0.7 | 3.6 | 0.4 | 4.1 |
| Iowa | 0.1 | (a) | 0.1 | 0.1 | 0.2 |
| Kansas | 0.1 | (a) | 0.1 | (a) | 0.1 |
| Kentucky | 13.2 | 6.2 | 19.4 | 1.3 | 20.7 |
| Missouri | 9.5 | 1.4 | 10.9 | 0.8 | 11.8 |
| Nebraska | 2.4 | 0.3 | 2.7 | 0.3 | 3.1 |
| Ohio | 4.8 | 2.4 | 7.2 | 0.2 | 7.4 |
| Total, Central States | 34.9 | 11.3 | 46.2 | 3.4 | 49.6 |
| Total, North | 770.5 | 242.3 | 1,012.8 | 78.0 | 1,090.8 |
| North Carolina | 219.0 | 24.4 | 243.4 | 1.9 | 245.3 |
| South Carolina | 175.0 | 18.6 | 193.6 | 2.4 | 196.0 |
| Virginia | 114.5 | 14.0 | 128.5 | 2.2 | 130.7 |
| Total, South Atlantic | 508.5 | 57.0 | 565.5 | 6.5 | 572.0 |
| Florida | 182.4 | 23.1 | 205.5 | 2.9 | 208.4 |
| Georgia | 307.8 | 34.7 | 342.5 | 1.9 | 344.4 |
| Total, East Gulf | 490.2 | 57.8 | 548.0 | 4.8 | 552.8 |

(continued)

Table 1.--Softwood tree biomass on timberland in the United States, by State and section, and class of timber, 1987, continued

| State and section | Live sound trees | | | Cull trees | Total tree biomass |
|--------------------------------|------------------|-----------|---------|---------------|-----------------------|
| | Tree boles | Tree tops | Total | | |
| ----- (Million dry tons) ----- | | | | | |
| Alabama | 214.5 | 41.6 | 256.1 | 17.6 | 273.7 |
| Mississippi | 168.4 | 31.0 | 199.4 | 8.6 | 208.0 |
| Tennessee | 50.8 | 9.4 | 60.2 | 4.4 | 64.6 |
| Total, Central Gulf | 433.7 | 82.0 | 515.7 | 30.6 | 546.3 |
| Arkansas | 137.6 | 23.9 | 161.5 | 5.9 | 167.4 |
| Louisiana | 193.4 | 34.3 | 227.7 | 9.3 | 237.0 |
| Oklahoma | 18.1 | 3.1 | 21.2 | 1.0 | 22.2 |
| Texas | 143.5 | 25.8 | 169.3 | 4.9 | 174.2 |
| Total, West Gulf | 492.6 | 87.1 | 579.7 | 21.1 | 600.8 |
| Total, South | 1,925.0 | 283.9 | 2,208.9 | 63.0 | 2,271.9 |
| Idaho | 400.8 | 100.2 | 501.0 | 51.2 | 552.2 |
| Montana | 349.8 | 116.6 | 466.4 | 153.4 | 619.8 |
| Wyoming | 75.8 | 26.6 | 102.5 | 11.1 | 113.6 |
| Total, N. Rocky Mountain | 826.5 | 243.5 | 1,069.9 | 215.7 | 1,285.6 |
| Arizona | 70.9 | 30.4 | 101.3 | 12.1 | 113.4 |
| Colorado | 181.1 | 57.2 | 238.3 | 30.8 | 269.1 |
| Nevada | 4.5 | 1.4 | 6.0 | 0.3 | 6.3 |
| New Mexico | 67.6 | 25.0 | 92.5 | 12.6 | 105.2 |
| Utah | 43.9 | 15.4 | 59.3 | 5.8 | 65.0 |
| Total, S. Rocky Mountain | 368.0 | 129.4 | 497.4 | 61.7 | 559.0 |
| California | 656.4 | 99.4 | 755.8 | 32.6 | 788.4 |
| Oregon | 882.8 | 150.7 | 1,033.5 | 12.8 | 1,046.3 |
| Washington | 799.1 | 150.3 | 949.4 | 5.2 | 954.6 |
| Total, Pacific Coast | 2,338.3 | 400.4 | 2,738.7 | 50.6 | 2,789.3 |
| Coastal Alaska | 414.8 | 75.6 | 490.4 | 23.4 | 513.8 |
| Interior Alaska | 108.6 | 36.0 | 144.6 | 1.3 | 145.9 |
| Hawaii | 0.1 | (a) | 0.1 | (a) | 0.1 |
| Total, Alaska and Hawaii | 523.5 | 111.6 | 635.1 | 24.7 | 659.8 |
| Total, West | 4,056.3 | 884.8 | 4,941.1 | 352.7 | 5,293.8 |
| Total, all States | 6,751.8 | 1,411.0 | 8,162.8 | 493.7 | 8,656.5 |

^a Less than 50,000 dry tons.

Table 2.--Hardwood tree biomass on timberland in the United States, by State and section, and class of timber, 1987

| State and section | Live sound trees | | | Cull trees | Total tree biomass |
|-------------------------|--------------------------------|-----------|---------|---------------|-----------------------|
| | Tree boles | Tree tops | Total | | |
| | ----- (Million dry tons) ----- | | | | |
| Connecticut | 40.6 | 14.6 | 55.2 | 3.7 | 58.9 |
| Maine | 154.7 | 59.3 | 214.0 | 52.0 | 266.1 |
| Massachusetts | 57.8 | 21.5 | 79.3 | 11.6 | 90.9 |
| New Hampshire | 78.9 | 29.7 | 108.5 | 17.0 | 125.6 |
| Rhode Island | 7.2 | 2.7 | 9.9 | 0.9 | 10.8 |
| Vermont | 74.0 | 27.2 | 101.2 | 24.7 | 125.9 |
| Total, New England | 413.3 | 154.9 | 568.2 | 110.0 | 678.2 |
| Delaware | 9.2 | 3.2 | 12.5 | 0.8 | 13.3 |
| Maryland | 71.6 | 24.6 | 96.2 | 5.4 | 101.6 |
| New Jersey | 21.3 | 7.7 | 29.0 | 2.8 | 31.8 |
| New York | 317.0 | 102.7 | 419.7 | 45.0 | 464.7 |
| Pennsylvania | 407.8 | 120.6 | 528.4 | 10.6 | 538.9 |
| West Virginia | 277.3 | 96.0 | 373.3 | 46.1 | 419.4 |
| Total, Middle Atlantic | 1,104.2 | 354.8 | 1,459.0 | 110.8 | 1,569.8 |
| Michigan | 251.2 | 78.8 | 330.0 | 24.9 | 354.9 |
| Minnesota | 155.2 | 46.5 | 201.7 | 32.4 | 234.1 |
| North Dakota | 3.9 | 1.2 | 5.1 | 2.0 | 7.0 |
| South Dakota | 1.1 | 0.3 | 1.4 | 1.2 | 2.6 |
| Wisconsin | 214.0 | 67.5 | 281.5 | 38.5 | 320.0 |
| Total, Lake States | 625.4 | 194.2 | 819.6 | 98.9 | 918.6 |
| Illinois | 89.9 | 26.0 | 116.0 | 18.9 | 134.8 |
| Indiana | 93.5 | 26.8 | 120.3 | 28.3 | 148.6 |
| Iowa | 22.6 | 6.7 | 29.3 | 14.0 | 43.3 |
| Kansas | 14.9 | 4.3 | 19.2 | 9.1 | 28.3 |
| Kentucky | 188.7 | 45.4 | 234.1 | 36.3 | 270.5 |
| Missouri | 148.6 | 48.8 | 197.5 | 96.8 | 294.3 |
| Nebraska | 5.0 | 1.3 | 6.2 | 2.8 | 9.0 |
| Ohio | 91.5 | 28.1 | 119.6 | 13.8 | 133.5 |
| Total, Central States | 654.7 | 187.4 | 842.2 | 220.1 | 1,062.2 |
| Total, North | 2,797.6 | 891.3 | 3,689.0 | 539.8 | 4,228.8 |
| North Carolina | 376.8 | 54.8 | 431.6 | 48.6 | 480.2 |
| South Carolina | 180.0 | 26.8 | 206.8 | 34.1 | 240.9 |
| Virginia | 371.5 | 57.0 | 428.5 | 60.9 | 489.4 |
| Total, South Atlantic | 928.3 | 138.6 | 1,066.9 | 143.6 | 1,210.5 |
| Florida | 110.9 | 16.5 | 127.4 | 46.3 | 173.7 |
| Georgia | 308.6 | 46.7 | 355.3 | 46.2 | 401.5 |
| Total, East Gulf | 419.5 | 63.2 | 482.7 | 92.5 | 575.2 |

(continued)

Table 2.--Hardwood tree biomass on timberland in the United States, by State and section, and class of timber, 1987, continued

| State and section | Live sound trees | | | Cull trees | Total tree biomass |
|--------------------------|--------------------------------|-----------|---------|---------------|-----------------------|
| | Tree boles | Tree tops | Total | | |
| | ----- (Million dry tons) ----- | | | | |
| Alabama | 240.5 | 80.3 | 320.8 | 106.2 | 427.0 |
| Mississippi | 242.6 | 86.5 | 329.1 | 102.1 | 431.2 |
| Tennessee | 288.2 | 94.2 | 382.4 | 131.9 | 514.3 |
| Total, Central Gulf | 771.3 | 261.0 | 1,032.3 | 340.2 | 1,372.5 |
| Arkansas | 255.3 | 94.5 | 349.8 | 96.9 | 446.7 |
| Louisiana | 195.0 | 69.0 | 264.0 | 104.9 | 368.9 |
| Oklahoma | 30.9 | 11.8 | 42.7 | 36.4 | 79.1 |
| Texas | 110.7 | 39.8 | 150.5 | 70.4 | 220.9 |
| Total, West Gulf | 591.9 | 215.1 | 807.0 | 308.6 | 1,115.6 |
| Total, South | 2,711.0 | 677.9 | 3,388.9 | 884.9 | 4,273.8 |
| Idaho | 5.2 | 1.1 | 6.3 | 1.4 | 7.7 |
| Montana | 4.2 | 1.3 | 5.6 | 0.7 | 6.3 |
| Wyoming | 3.5 | 0.9 | 4.4 | 1.5 | 5.9 |
| Total, N. Rocky Mountain | 13.0 | 3.3 | 16.2 | 3.6 | 19.8 |
| Arizona | 3.5 | 0.8 | 4.3 | 3.6 | 7.8 |
| Colorado | 33.2 | 6.8 | 40.0 | 10.6 | 50.6 |
| Nevada | 0.3 | 0.1 | 0.4 | (a) | 0.4 |
| New Mexico | 5.1 | 1.0 | 6.2 | 0.9 | 7.0 |
| Utah | 9.1 | 2.0 | 11.1 | 0.8 | 11.9 |
| Total, S. Rocky Mountain | 51.1 | 10.7 | 61.8 | 15.9 | 77.7 |
| California | 165.2 | 37.4 | 202.6 | 27.4 | 230.0 |
| Oregon | 80.4 | 14.5 | 94.9 | 10.1 | 105.0 |
| Washington | 96.8 | 19.4 | 116.2 | 6.7 | 122.9 |
| Total, Pacific Coast | 342.5 | 71.3 | 413.8 | 44.2 | 458.0 |
| Coastal Alaska | 3.0 | 1.0 | 4.0 | 0.1 | 4.1 |
| Interior Alaska | 81.0 | 17.0 | 98.1 | 8.2 | 106.3 |
| Hawaii | 6.2 | 1.5 | 7.7 | 1.1 | 8.9 |
| Total, Alaska and Hawaii | 90.3 | 19.5 | 109.8 | 9.5 | 119.3 |
| Total, West | 496.8 | 104.8 | 601.7 | 73.1 | 674.8 |
| Total, all States | 6,005.4 | 1,674.1 | 7,679.5 | 1,497.9 | 9,177.4 |

^a Less than 50,000 dry tons.

Table 3.--Total tree biomass on timberland in the United States, by State and section, and class of timber, 1987

| State and section | Live sound trees | | | Cull trees | Total tree biomass |
|-------------------------|--------------------------------|-----------|---------|---------------|-----------------------|
| | Tree boles | Tree tops | Total | | |
| | ----- (Million dry tons) ----- | | | | |
| Connecticut | 47.8 | 17.0 | 64.8 | 4.2 | 69.0 |
| Maine | 374.5 | 144.6 | 519.2 | 82.6 | 601.8 |
| Massachusetts | 86.3 | 30.8 | 117.2 | 13.8 | 130.9 |
| New Hampshire | 135.2 | 49.8 | 185.0 | 23.9 | 208.9 |
| Rhode Island | 8.0 | 3.0 | 11.0 | 0.9 | 12.0 |
| Vermont | 105.4 | 38.7 | 144.0 | 32.5 | 176.6 |
| Total, New England | 757.2 | 283.9 | 1,041.2 | 157.9 | 1,199.1 |
| Delaware | 11.8 | 4.2 | 15.9 | 0.8 | 16.7 |
| Maryland | 83.9 | 29.1 | 113.0 | 5.6 | 118.6 |
| New Jersey | 30.9 | 11.5 | 42.3 | 4.8 | 47.2 |
| New York | 418.4 | 135.9 | 554.4 | 55.1 | 609.5 |
| Pennsylvania | 440.0 | 130.6 | 570.6 | 11.0 | 581.5 |
| West Virginia | 293.5 | 101.8 | 395.3 | 48.0 | 443.2 |
| Total, Middle Atlantic | 1,278.5 | 413.0 | 1,691.5 | 125.2 | 1,816.7 |
| Michigan | 336.0 | 97.0 | 432.9 | 29.7 | 462.6 |
| Minnesota | 209.1 | 58.6 | 267.7 | 36.9 | 304.6 |
| North Dakota | 4.0 | 1.2 | 5.1 | 2.0 | 7.1 |
| South Dakota | 24.7 | 3.4 | 28.1 | 1.5 | 29.5 |
| Wisconsin | 269.1 | 77.9 | 347.0 | 41.1 | 388.1 |
| Total, Lake States | 842.8 | 238.0 | 1,080.8 | 111.2 | 1,192.0 |
| Illinois | 91.7 | 26.3 | 118.0 | 19.0 | 137.0 |
| Indiana | 96.5 | 27.4 | 123.9 | 28.8 | 152.7 |
| Iowa | 22.7 | 6.7 | 29.4 | 14.1 | 43.5 |
| Kansas | 15.0 | 4.3 | 19.3 | 9.1 | 28.5 |
| Kentucky | 201.9 | 51.6 | 253.5 | 37.7 | 291.2 |
| Missouri | 158.2 | 50.2 | 208.4 | 97.7 | 306.1 |
| Nebraska | 7.4 | 1.6 | 9.0 | 3.1 | 12.0 |
| Ohio | 96.3 | 30.6 | 126.8 | 14.0 | 140.8 |
| Total, Central States | 689.6 | 198.7 | 888.4 | 223.5 | 1,111.8 |
| Total, North | 3,568.2 | 1,133.6 | 4,701.8 | 617.8 | 5,319.6 |
| North Carolina | 595.8 | 79.2 | 675.0 | 50.5 | 725.5 |
| South Carolina | 355.0 | 45.4 | 400.4 | 36.5 | 436.9 |
| Virginia | 486.0 | 71.0 | 557.0 | 63.1 | 620.1 |
| Total, South Atlantic | 1,436.8 | 195.6 | 1,632.4 | 150.1 | 1,782.5 |
| Florida | 293.3 | 39.6 | 332.9 | 49.2 | 382.1 |
| Georgia | 616.4 | 81.4 | 697.8 | 48.1 | 745.9 |
| Total, East Gulf | 909.7 | 121.0 | 1,030.7 | 97.3 | 1,128.0 |

(continued)

Table 3.--Total tree biomass on timberland in the United States, by State and section, and class of timber, 1987, continued

| State and section | Live sound trees | | | Cull trees | Total tree biomass |
|--------------------------|--------------------------------|-----------|----------|---------------|-----------------------|
| | Tree boles | Tree tops | Total | | |
| | ----- (Million dry tons) ----- | | | | |
| Alabama | 455.0 | 121.9 | 576.9 | 123.8 | 700.7 |
| Mississippi | 411.0 | 117.5 | 528.5 | 110.7 | 639.2 |
| Tennessee | 339.0 | 103.6 | 442.6 | 136.3 | 578.9 |
| Total, Central Gulf | 1,205.0 | 343.0 | 1,548.0 | 370.8 | 1,918.8 |
| Arkansas | 392.9 | 118.4 | 511.3 | 102.8 | 614.1 |
| Louisiana | 388.4 | 103.3 | 491.7 | 114.2 | 605.9 |
| Oklahoma | 49.0 | 14.9 | 63.9 | 37.4 | 101.3 |
| Texas | 254.2 | 65.6 | 319.8 | 75.3 | 395.1 |
| Total, West Gulf | 1,084.5 | 302.2 | 1,386.7 | 329.7 | 1,716.4 |
| Total, South | 4,636.0 | 961.8 | 5,597.8 | 947.9 | 6,545.7 |
| Idaho | 406.0 | 101.3 | 507.3 | 52.6 | 559.9 |
| Montana | 354.1 | 118.0 | 472.0 | 154.1 | 626.1 |
| Wyoming | 79.3 | 27.5 | 106.9 | 12.6 | 119.4 |
| Total, N. Rocky Mountain | 839.4 | 246.7 | 1,086.1 | 219.3 | 1,305.5 |
| Arizona | 74.4 | 31.2 | 105.6 | 15.7 | 121.3 |
| Colorado | 214.3 | 64.0 | 278.2 | 41.4 | 319.7 |
| Nevada | 4.8 | 1.5 | 6.3 | 0.3 | 6.7 |
| New Mexico | 72.7 | 26.0 | 98.7 | 13.5 | 112.2 |
| Utah | 52.9 | 17.4 | 70.3 | 6.6 | 76.9 |
| Total, S. Rocky Mountain | 419.1 | 140.1 | 559.2 | 77.5 | 636.7 |
| California | 821.6 | 136.8 | 958.4 | 59.9 | 1,018.4 |
| Oregon | 963.2 | 165.2 | 1,128.4 | 22.9 | 1,151.3 |
| Washington | 896.0 | 169.7 | 1,065.6 | 12.0 | 1,077.6 |
| Total, Pacific Coast | 2,680.8 | 471.7 | 3,152.5 | 94.7 | 3,247.2 |
| Coastal Alaska | 417.9 | 76.5 | 494.4 | 23.5 | 518.0 |
| Interior Alaska | 189.7 | 53.0 | 242.7 | 9.5 | 252.2 |
| Hawaii | 6.3 | 1.5 | 7.8 | 1.1 | 9.0 |
| Total, Alaska and Hawaii | 613.8 | 131.1 | 744.9 | 34.2 | 779.1 |
| Total, West | 4,553.1 | 989.7 | 5,542.8 | 425.8 | 5,968.6 |
| Total, all States | 12,757.3 | 3,085.1 | 15,842.4 | 1,991.6 | 17,833.9 |

^a Less than 50,000 dry tons.

Table 4.--Softwood tree biomass on timberland in the United States, by State and section, and ownership class, 1987

| State and section | Public lands | | | Private lands | | | All ownerships |
|-------------------------|--------------------------------|-----------------|-----------------|--------------------|------------------|------------------|-------------------|
| | National Forest | Other public | Total public | Forest industry | Other private | Total private | |
| | ----- (Million dry tons) ----- | | | | | | |
| Connecticut | 0.0 | 1.4 | 1.4 | 0.0 | 8.7 | 8.7 | 10.1 |
| Maine | 0.9 | 11.1 | 12.0 | 162.0 | 161.7 | 323.7 | 335.7 |
| Massachusetts | 0.0 | 6.3 | 6.3 | 1.1 | 32.7 | 33.8 | 40.1 |
| New Hampshire | 8.8 | 4.9 | 13.7 | 11.5 | 58.1 | 69.6 | 83.3 |
| Rhode Island | 0.0 | 0.3 | 0.3 | 0.0 | 0.9 | 0.9 | 1.1 |
| Vermont | 2.9 | 4.7 | 7.6 | 4.0 | 39.1 | 43.1 | 50.7 |
| Total, New England | 12.5 | 28.6 | 41.2 | 178.5 | 301.2 | 479.7 | 520.9 |
| Delaware | 0.0 | 0.1 | 0.1 | 0.3 | 3.1 | 3.3 | 3.4 |
| Maryland | 0.0 | 1.9 | 1.9 | 0.9 | 14.1 | 15.1 | 17.0 |
| New Jersey | 0.0 | 4.3 | 4.3 | 0.0 | 11.0 | 11.0 | 15.3 |
| New York | 0.1 | 11.1 | 11.1 | 10.2 | 123.4 | 133.7 | 144.8 |
| Pennsylvania | 1.3 | 8.0 | 9.3 | 2.4 | 30.9 | 33.3 | 42.6 |
| West Virginia | 1.8 | 0.8 | 2.7 | 2.1 | 19.0 | 21.1 | 23.8 |
| Total, Middle Atlantic | 3.2 | 26.3 | 29.5 | 15.9 | 201.6 | 217.5 | 246.9 |
| Michigan | 15.4 | 23.8 | 39.2 | 12.2 | 56.4 | 68.6 | 107.8 |
| Minnesota | 8.7 | 29.2 | 37.8 | 4.1 | 28.6 | 32.7 | 70.6 |
| North Dakota | 0.0 | (a) | (a) | 0.0 | (a) | (a) | (a) |
| South Dakota | 17.0 | 2.4 | 19.4 | 0.4 | 7.1 | 7.5 | 26.9 |
| Wisconsin | 5.7 | 15.2 | 20.9 | 5.4 | 41.8 | 47.2 | 68.1 |
| Total, Lake States | 46.8 | 70.6 | 117.4 | 22.1 | 134.0 | 156.0 | 273.4 |
| Illinois | 0.1 | 0.1 | 0.2 | (a) | 1.9 | 1.9 | 2.2 |
| Indiana | 0.2 | 0.4 | 0.5 | (a) | 3.5 | 3.5 | 4.1 |
| Iowa | 0.0 | (a) | (a) | 0.0 | 0.2 | 0.2 | 0.2 |
| Kansas | 0.0 | (a) | (a) | 0.0 | 0.2 | 0.2 | 0.1 |
| Kentucky | 1.0 | 0.5 | 1.6 | 0.4 | 18.8 | 19.2 | 20.7 |
| Missouri | 1.3 | 0.3 | 1.6 | 0.2 | 9.9 | 10.1 | 11.8 |
| Nebraska | 0.2 | 0.2 | 0.4 | 0.0 | 2.7 | 2.7 | 3.1 |
| Ohio | 0.2 | 0.3 | 0.4 | 0.1 | 6.8 | 7.0 | 7.4 |
| Total, Central States | 2.9 | 1.8 | 4.7 | 0.7 | 44.2 | 44.9 | 49.6 |
| Total, North | 65.5 | 127.2 | 192.7 | 217.1 | 681.0 | 898.1 | 1,090.8 |
| North Carolina | 10.2 | 11.7 | 21.9 | 33.7 | 189.7 | 223.4 | 245.3 |
| South Carolina | 15.9 | 13.1 | 29.0 | 40.1 | 126.9 | 167.0 | 196.0 |
| Virginia | 6.4 | 7.3 | 13.7 | 24.3 | 92.7 | 117.0 | 130.7 |
| Total, South Atlantic | 32.5 | 32.1 | 64.6 | 98.1 | 409.3 | 507.4 | 572.0 |
| Florida | 19.4 | 26.2 | 45.6 | 61.0 | 101.8 | 162.8 | 208.4 |
| Georgia | 9.4 | 21.5 | 30.9 | 76.1 | 237.4 | 313.5 | 344.4 |
| Total, East Gulf | 28.8 | 47.7 | 76.5 | 137.1 | 339.2 | 476.3 | 552.8 |

(continued)

Table 4.--Softwood tree biomass on timberland in the United States, by State and section, and ownership class, 1987, continued

| State and section | Public lands | | | Private lands | | | All ownerships |
|--------------------------|--------------------|-----------------|-----------------|--------------------|------------------|------------------|-------------------|
| | National Forest | Other public | Total public | Forest industry | Other private | Total private | |
| | (Million dry tons) | | | | | | |
| Alabama | 14.8 | 5.4 | 20.2 | 66.2 | 187.3 | 253.5 | 273.7 |
| Mississippi | 29.0 | 9.8 | 38.8 | 39.3 | 129.9 | 169.2 | 208.0 |
| Tennessee | 6.7 | 5.3 | 12.0 | 6.5 | 46.1 | 52.6 | 64.6 |
| Total, Central Gulf | 50.5 | 20.5 | 71.0 | 112.0 | 363.3 | 475.3 | 546.3 |
| Arkansas | 31.6 | 4.5 | 36.1 | 59.9 | 71.4 | 131.3 | 167.4 |
| Louisiana | 17.3 | 6.0 | 23.3 | 62.9 | 150.8 | 213.7 | 237.0 |
| Oklahoma | 3.5 | 1.3 | 4.8 | 7.7 | 9.7 | 17.4 | 22.2 |
| Texas | 25.4 | 2.8 | 28.2 | 50.3 | 95.7 | 146.0 | 174.2 |
| Total, West Gulf | 77.8 | 14.6 | 92.4 | 180.8 | 327.6 | 508.4 | 600.8 |
| Total, South | 189.6 | 114.9 | 304.5 | 528.0 | 1,439.4 | 1,967.4 | 2,271.9 |
| Idaho | 368.8 | 65.7 | 434.4 | 45.5 | 72.2 | 117.8 | 552.2 |
| Montana | 349.1 | 71.6 | 420.8 | 71.6 | 127.5 | 199.1 | 619.8 |
| Wyoming | 58.0 | 20.7 | 78.6 | 1.0 | 34.0 | 34.9 | 113.6 |
| Total, N. Rocky Mountain | 775.8 | 158.0 | 933.9 | 118.1 | 233.7 | 351.8 | 1,285.6 |
| Arizona | 74.0 | 38.2 | 112.2 | 0.0 | 1.3 | 1.3 | 113.4 |
| Colorado | 161.9 | 33.3 | 195.1 | 0.0 | 73.9 | 73.9 | 269.1 |
| Nevada | 2.8 | 0.3 | 3.1 | 0.0 | 3.2 | 3.2 | 6.3 |
| New Mexico | 58.1 | 14.7 | 72.8 | 0.1 | 32.3 | 32.4 | 105.2 |
| Utah | 44.5 | 8.5 | 53.0 | 0.0 | 12.0 | 12.0 | 65.0 |
| Total, S. Rocky Mountain | 341.3 | 94.9 | 436.2 | 0.1 | 122.7 | 122.8 | 559.0 |
| California | 412.4 | 24.3 | 436.7 | 130.0 | 221.6 | 351.7 | 788.4 |
| Oregon | 488.4 | 154.8 | 643.2 | 246.0 | 157.1 | 403.1 | 1,046.3 |
| Washington | 275.3 | 214.9 | 490.2 | 260.0 | 204.5 | 464.5 | 954.6 |
| Total, Pacific Coast | 1,176.1 | 394.0 | 1,570.1 | 636.0 | 583.2 | 1,219.2 | 2,789.3 |
| Coastal Alaska | 403.6 | 47.7 | 451.3 | 0.0 | 62.6 | 62.6 | 513.8 |
| Interior Alaska | 0.0 | 66.6 | 66.6 | 0.0 | 79.3 | 79.3 | 145.9 |
| Hawaii | 0.0 | (a) | (a) | 0.0 | (a) | (a) | 0.1 |
| Total, Alaska and Hawaii | 403.6 | 114.3 | 517.9 | 0.0 | 141.9 | 141.9 | 659.8 |
| Total, West | 2,696.8 | 761.2 | 3,458.0 | 754.3 | 1,081.4 | 1,835.7 | 5,293.8 |
| Total, all States | 2,951.9 | 1,003.4 | 3,955.3 | 1,499.5 | 3,201.8 | 4,701.3 | 8,656.5 |

^a Less than 50,000 dry tons.

Table 5.--Hardwood tree biomass on timberland in the United States, by State and section, and ownership class, 1987

| State and section | Public lands | | | Private lands | | | All ownerships |
|-------------------------|--------------------------------|-----------------|-----------------|--------------------|------------------|------------------|-------------------|
| | National Forest | Other public | Total public | Forest industry | Other private | Total private | |
| | ----- (Million dry tons) ----- | | | | | | |
| Connecticut | 0.0 | 8.2 | 8.2 | 0.0 | 50.7 | 50.7 | 58.9 |
| Maine | 0.7 | 8.8 | 9.5 | 128.4 | 128.2 | 256.6 | 266.1 |
| Massachusetts | 0.0 | 14.3 | 14.3 | 2.4 | 74.1 | 76.6 | 90.9 |
| New Hampshire | 13.2 | 7.4 | 20.6 | 17.3 | 87.7 | 105.0 | 125.6 |
| Rhode Island | 0.0 | 2.4 | 2.4 | 0.0 | 8.4 | 8.4 | 10.8 |
| Vermont | 7.1 | 11.6 | 18.8 | 10.0 | 97.1 | 107.1 | 125.9 |
| Total, New England | 21.1 | 52.7 | 73.8 | 158.2 | 446.3 | 604.4 | 678.2 |
| Delaware | 0.0 | 0.5 | 0.5 | 1.0 | 11.8 | 12.8 | 13.3 |
| Maryland | 0.0 | 11.5 | 11.5 | 5.5 | 84.6 | 90.1 | 101.6 |
| New Jersey | 0.0 | 8.9 | 8.9 | 0.0 | 23.0 | 23.0 | 31.8 |
| New York | 0.2 | 35.6 | 35.8 | 32.8 | 396.1 | 428.9 | 464.7 |
| Pennsylvania | 16.2 | 101.8 | 118.0 | 29.8 | 391.1 | 420.9 | 538.9 |
| West Virginia | 32.6 | 14.4 | 47.0 | 36.8 | 335.6 | 372.5 | 419.4 |
| Total, Middle Atlantic | 49.0 | 172.7 | 221.6 | 105.9 | 1,242.2 | 1,348.1 | 1,569.8 |
| Michigan | 50.6 | 78.4 | 129.0 | 40.2 | 185.7 | 225.9 | 354.9 |
| Minnesota | 28.8 | 96.7 | 125.5 | 13.6 | 95.0 | 108.6 | 234.1 |
| North Dakota | 0.0 | 1.4 | 1.4 | 0.0 | 5.7 | 5.7 | 7.0 |
| South Dakota | 1.7 | 0.2 | 1.9 | (a) | 0.7 | 0.7 | 2.6 |
| Wisconsin | 27.0 | 71.3 | 98.3 | 25.2 | 196.6 | 221.7 | 320.0 |
| Total, Lake States | 108.0 | 248.0 | 356.0 | 79.0 | 483.6 | 562.5 | 918.6 |
| Illinois | 7.6 | 5.5 | 13.0 | 0.4 | 121.4 | 121.8 | 134.8 |
| Indiana | 5.7 | 12.8 | 18.5 | 0.6 | 129.5 | 130.1 | 148.6 |
| Iowa | 0.0 | 3.0 | 3.0 | 0.0 | 40.2 | 40.2 | 43.3 |
| Kansas | 0.0 | 1.2 | 1.2 | 0.0 | 27.1 | 27.1 | 28.3 |
| Kentucky | 13.2 | 7.0 | 20.2 | 4.7 | 245.6 | 250.2 | 270.5 |
| Missouri | 32.0 | 8.7 | 40.7 | 5.7 | 248.0 | 253.7 | 294.3 |
| Nebraska | 0.5 | 0.6 | 1.1 | 0.0 | 7.9 | 7.9 | 9.0 |
| Ohio | 3.2 | 4.7 | 7.8 | 2.1 | 123.6 | 125.6 | 133.5 |
| Total, Central States | 62.2 | 43.3 | 105.5 | 13.4 | 943.3 | 956.8 | 1,062.2 |
| Total, North | 240.2 | 516.6 | 756.9 | 356.5 | 3,115.4 | 3,471.9 | 4,228.8 |
| North Carolina | 49.4 | 14.1 | 63.5 | 36.7 | 380.0 | 416.7 | 480.2 |
| South Carolina | 10.3 | 9.3 | 19.6 | 42.0 | 179.3 | 221.3 | 240.9 |
| Virginia | 52.1 | 20.8 | 72.9 | 29.9 | 386.6 | 416.5 | 489.4 |
| Total, South Atlantic | 111.8 | 44.2 | 156.0 | 108.6 | 945.9 | 1,054.5 | 1,210.5 |
| Florida | 6.2 | 24.2 | 30.4 | 41.2 | 102.1 | 143.3 | 173.7 |
| Georgia | 25.0 | 18.1 | 43.1 | 51.6 | 306.8 | 358.4 | 401.5 |
| Total, East Gulf | 31.2 | 42.3 | 73.5 | 92.8 | 408.9 | 501.7 | 575.2 |

(continued)

Table 5.--Hardwood tree biomass on timberland in the United States, by State and section, and ownership class, 1987, continued

| State and section | Public lands | | | Private lands | | | All ownerships |
|--------------------------|--------------------|-----------------|-----------------|--------------------|------------------|------------------|-------------------|
| | National Forest | Other public | Total public | Forest industry | Other private | Total private | |
| | (Million dry tons) | | | | | | |
| Alabama | 13.0 | 13.9 | 26.9 | 69.5 | 330.6 | 400.1 | 427.0 |
| Mississippi | 29.0 | 24.9 | 53.9 | 52.9 | 324.4 | 377.3 | 431.2 |
| Tennessee | 26.8 | 31.7 | 58.5 | 41.4 | 414.4 | 455.8 | 514.3 |
| Total, Central Gulf | 68.8 | 70.5 | 139.3 | 163.8 | 1,069.4 | 1,233.2 | 1,372.5 |
| Arkansas | 62.0 | 35.2 | 97.2 | 77.0 | 272.5 | 349.5 | 446.7 |
| Louisiana | 12.1 | 28.8 | 40.9 | 68.8 | 259.2 | 328.0 | 368.9 |
| Oklahoma | 4.4 | 7.2 | 11.6 | 8.3 | 59.2 | 67.5 | 79.1 |
| Texas | 8.5 | 4.3 | 12.8 | 54.5 | 153.6 | 208.1 | 220.9 |
| Total, West Gulf | 87.0 | 75.5 | 162.5 | 208.6 | 744.5 | 953.1 | 1,115.6 |
| Total, South | 298.8 | 232.5 | 531.3 | 573.8 | 3,168.7 | 3,742.5 | 4,273.8 |
| Idaho | 5.1 | 0.9 | 6.1 | 0.6 | 1.0 | 1.6 | 7.7 |
| Montana | 3.5 | 0.7 | 4.2 | 0.7 | 1.3 | 2.0 | 6.3 |
| Wyoming | 3.0 | 1.1 | 4.1 | (a) | 1.8 | 1.8 | 5.9 |
| Total, N. Rocky Mountain | 11.7 | 2.7 | 14.4 | 1.4 | 4.0 | 5.4 | 19.8 |
| Arizona | 5.1 | 2.6 | 7.7 | 0.0 | 0.1 | 0.1 | 7.8 |
| Colorado | 30.4 | 6.3 | 36.7 | 0.0 | 13.9 | 13.9 | 50.6 |
| Nevada | 0.2 | (a) | 0.2 | 0.0 | 0.2 | 0.2 | 0.4 |
| New Mexico | 3.9 | 1.0 | 4.9 | (a) | 2.2 | 2.2 | 7.0 |
| Utah | 8.1 | 1.6 | 9.7 | 0.0 | 2.2 | 2.2 | 11.9 |
| Total, S. Rocky Mountain | 47.7 | 11.4 | 59.2 | (a) | 18.5 | 18.5 | 77.7 |
| California | 120.3 | 7.1 | 127.4 | 37.9 | 64.7 | 102.6 | 230.0 |
| Oregon | 49.0 | 15.5 | 64.6 | 24.7 | 15.8 | 40.5 | 105.0 |
| Washington | 35.5 | 27.7 | 63.1 | 33.5 | 26.3 | 59.8 | 122.9 |
| Total, Pacific Coast | 204.8 | 50.3 | 255.1 | 96.1 | 106.8 | 202.9 | 458.0 |
| Coastal Alaska | 3.2 | 0.4 | 3.6 | 0.0 | 0.5 | 0.5 | 4.1 |
| Interior Alaska | 0.0 | 48.5 | 48.5 | 0.0 | 57.8 | 57.8 | 106.3 |
| Hawaii | 0.0 | 4.3 | 4.3 | 0.0 | 4.6 | 4.6 | 8.9 |
| Total, Alaska and Hawaii | 3.2 | 53.2 | 56.4 | 0.0 | 62.9 | 62.9 | 119.3 |
| Total, West | 267.4 | 117.6 | 385.1 | 97.5 | 192.2 | 289.7 | 674.8 |
| Total, all States | 806.5 | 866.8 | 1,673.3 | 1,027.9 | 6,476.3 | 7,504.1 | 9,177.4 |

^a Less than 50,000 dry tons.

Table 6.--Total tree biomass on timberland in the United States, by State and section, and ownership class, 1987

| State and section | Public lands | | | Private lands | | | All ownerships |
|-------------------------|--------------------------------|-----------------|-----------------|--------------------|------------------|------------------|-------------------|
| | National Forest | Other public | Total public | Forest industry | Other private | Total private | |
| | ----- (Million dry tons) ----- | | | | | | |
| Connecticut | 0.0 | 9.6 | 9.6 | 0.0 | 59.4 | 59.4 | 69.0 |
| Maine | 1.6 | 19.9 | 21.5 | 290.3 | 290.0 | 580.3 | 601.8 |
| Massachusetts | 0.0 | 20.6 | 20.6 | 3.5 | 106.8 | 110.3 | 130.9 |
| New Hampshire | 22.0 | 12.3 | 34.3 | 28.8 | 145.8 | 174.6 | 208.9 |
| Rhode Island | 0.0 | 2.7 | 2.7 | 0.0 | 9.3 | 9.3 | 12.0 |
| Vermont | 10.0 | 16.3 | 26.3 | 14.1 | 136.2 | 150.3 | 176.6 |
| Total, New England | 33.6 | 81.3 | 114.9 | 336.7 | 747.4 | 1,084.1 | 1,199.1 |
| Delaware | 0.0 | 0.6 | 0.6 | 1.3 | 14.8 | 16.1 | 16.7 |
| Maryland | 0.0 | 13.5 | 13.5 | 6.4 | 98.7 | 105.1 | 118.6 |
| New Jersey | 0.0 | 13.1 | 13.1 | 0.0 | 34.0 | 34.0 | 47.2 |
| New York | 0.2 | 46.7 | 46.9 | 43.1 | 519.5 | 562.6 | 609.5 |
| Pennsylvania | 17.5 | 109.9 | 127.4 | 32.1 | 422.0 | 454.2 | 581.5 |
| West Virginia | 34.4 | 15.2 | 49.6 | 38.9 | 354.7 | 393.6 | 443.2 |
| Total, Middle Atlantic | 52.1 | 198.9 | 251.1 | 121.8 | 1,443.8 | 1,565.6 | 1,816.7 |
| Michigan | 65.9 | 102.2 | 168.1 | 52.4 | 242.1 | 294.5 | 462.6 |
| Minnesota | 37.5 | 125.9 | 163.3 | 17.7 | 123.6 | 141.3 | 304.6 |
| North Dakota | 0.0 | 1.4 | 1.4 | 0.0 | 5.7 | 5.7 | 7.1 |
| South Dakota | 18.7 | 2.7 | 21.3 | 0.4 | 7.8 | 8.2 | 29.5 |
| Wisconsin | 32.7 | 86.5 | 119.2 | 30.5 | 238.4 | 268.9 | 388.1 |
| Total, Lake States | 154.8 | 318.5 | 473.4 | 101.0 | 617.6 | 718.7 | 1,192.0 |
| Illinois | 7.7 | 5.5 | 13.2 | 0.4 | 123.3 | 123.8 | 137.0 |
| Indiana | 5.9 | 13.1 | 19.0 | 0.6 | 133.0 | 133.7 | 152.7 |
| Iowa | 0.0 | 3.0 | 3.0 | 0.0 | 40.5 | 40.5 | 43.5 |
| Kansas | 0.0 | 1.2 | 1.2 | 0.0 | 27.3 | 27.3 | 28.5 |
| Kentucky | 14.3 | 7.5 | 21.8 | 5.0 | 264.4 | 269.4 | 291.2 |
| Missouri | 33.2 | 9.0 | 42.3 | 5.9 | 257.9 | 263.8 | 306.1 |
| Nebraska | 0.7 | 0.8 | 1.4 | 0.0 | 10.6 | 10.6 | 12.0 |
| Ohio | 3.3 | 4.9 | 8.2 | 2.2 | 130.4 | 132.6 | 140.8 |
| Total, Central States | 65.1 | 45.1 | 110.2 | 14.2 | 987.5 | 1,001.7 | 1,111.8 |
| Total, North | 305.7 | 643.9 | 949.6 | 573.7 | 3,796.4 | 4,370.1 | 5,319.6 |
| North Carolina | 59.6 | 25.8 | 85.4 | 70.4 | 569.7 | 640.1 | 725.5 |
| South Carolina | 26.2 | 22.4 | 48.6 | 82.1 | 306.2 | 388.3 | 436.9 |
| Virginia | 58.5 | 28.1 | 86.6 | 54.2 | 479.3 | 533.5 | 620.1 |
| Total, South Atlantic | 144.3 | 76.3 | 220.6 | 206.7 | 1,355.2 | 1,561.9 | 1,782.5 |
| Florida | 25.6 | 50.4 | 76.0 | 102.2 | 203.9 | 306.1 | 382.1 |
| Georgia | 34.4 | 39.6 | 74.0 | 127.7 | 544.2 | 671.9 | 745.9 |
| Total, East Gulf | 60.0 | 90.0 | 150.0 | 229.9 | 748.1 | 978.0 | 1,128.0 |

(continued)

Table 6.--Total tree biomass on timberland in the United States, by State and section, and ownership class, 1987, continued

| State and section | Public lands | | | Private lands | | | All ownerships |
|--------------------------|--------------------|-----------------|-----------------|--------------------|------------------|------------------|-------------------|
| | National Forest | Other public | Total public | Forest industry | Other private | Total private | |
| | (Million dry tons) | | | | | | |
| Alabama | 27.8 | 19.3 | 47.1 | 135.7 | 517.9 | 653.6 | 700.7 |
| Mississippi | 58.0 | 34.7 | 92.7 | 92.2 | 454.3 | 546.5 | 639.2 |
| Tennessee | 33.5 | 37.0 | 70.5 | 47.9 | 460.5 | 508.4 | 578.9 |
| Total, Central Gulf | 119.3 | 91.0 | 210.3 | 275.8 | 1,432.7 | 1,708.5 | 1,918.8 |
| Arkansas | 93.6 | 39.7 | 133.3 | 136.9 | 343.9 | 480.8 | 614.1 |
| Louisiana | 29.4 | 34.8 | 64.2 | 131.7 | 410.0 | 541.7 | 605.9 |
| Oklahoma | 7.9 | 8.5 | 16.4 | 16.0 | 68.9 | 84.9 | 101.3 |
| Texas | 33.9 | 7.1 | 41.0 | 104.8 | 249.3 | 354.1 | 395.1 |
| Total, West Gulf | 164.8 | 90.1 | 254.9 | 389.4 | 1,072.1 | 1,461.5 | 1,716.4 |
| Total, South | 488.4 | 347.4 | 835.8 | 1,101.8 | 4,608.1 | 5,709.9 | 13,091.4 |
| Idaho | 373.9 | 66.6 | 440.5 | 46.2 | 73.2 | 119.4 | 559.9 |
| Montana | 352.7 | 72.4 | 425.0 | 72.4 | 128.7 | 201.1 | 626.1 |
| Wyoming | 61.0 | 21.8 | 82.7 | 1.0 | 35.7 | 36.7 | 119.4 |
| Total, N. Rocky Mountain | 787.5 | 160.7 | 948.2 | 119.5 | 237.7 | 357.2 | 1,305.5 |
| Arizona | 79.1 | 40.8 | 119.9 | 0.0 | 1.4 | 1.4 | 121.3 |
| Colorado | 192.3 | 39.5 | 231.8 | 0.0 | 87.8 | 87.8 | 319.7 |
| Nevada | 3.0 | 0.3 | 3.3 | 0.0 | 3.4 | 3.4 | 6.7 |
| New Mexico | 62.0 | 15.7 | 77.7 | 0.1 | 34.4 | 34.5 | 112.2 |
| Utah | 52.7 | 10.1 | 62.7 | 0.0 | 14.2 | 14.2 | 76.9 |
| Total, S. Rocky Mountain | 389.1 | 106.4 | 495.4 | 0.1 | 141.2 | 141.3 | 636.7 |
| California | 532.7 | 31.4 | 564.1 | 168.0 | 286.3 | 454.3 | 1,018.4 |
| Oregon | 537.4 | 170.3 | 707.7 | 270.7 | 172.8 | 443.6 | 1,151.3 |
| Washington | 310.8 | 242.5 | 553.3 | 293.4 | 230.8 | 524.3 | 1,077.6 |
| Total, Pacific Coast | 1,380.9 | 444.3 | 1,825.1 | 732.2 | 689.9 | 1,422.1 | 3,247.2 |
| Coastal Alaska | 406.8 | 48.1 | 454.9 | 0.0 | 63.1 | 63.1 | 518.0 |
| Interior Alaska | 0.0 | 115.2 | 115.2 | 0.0 | 137.1 | 137.1 | 252.2 |
| Hawaii | 0.0 | 4.3 | 4.3 | 0.0 | 4.6 | 4.6 | 9.0 |
| Total, Alaska and Hawaii | 406.8 | 167.6 | 574.4 | 0.0 | 204.8 | 204.8 | 779.1 |
| Total, West | 2,964.2 | 878.9 | 3,843.2 | 851.8 | 1,273.6 | 2,125.4 | 5,968.6 |
| Total, all States | 3,758.3 | 1,870.2 | 5,628.5 | 2,527.3 | 9,678.1 | 12,205.4 | 17,833.9 |

^a Less than 50,000 dry tons.

Table 7.--Total forest biomass on timberland in the United States, by major species group, section, region, and class of material, 1987

| Species group, section, and region | Biomass of live sound and cull trees | Other live tree biomass | | Other forest biomass | | | Total biomass |
|--|--|-------------------------|---------|----------------------|----------|--------|------------------|
| | | Stumps | Foliage | Seedlings | Saplings | Shrubs | |
| ----- (Million dry tons) ----- | | | | | | | |
| Softwoods | 520.9 | 8.1 | (a) | (a) | 78.5 | (a) | 607.5 |
| Hardwoods | 678.2 | 12.1 | (a) | (a) | 102.6 | (a) | 792.9 |
| Total, New England | 1,199.1 | 20.2 | (a) | (a) | 181.1 | (a) | 1,400.4 |
| Softwoods | 246.9 | 2.9 | (a) | (a) | 31.6 | (a) | 281.5 |
| Hardwoods | 1,569.8 | 21.6 | (a) | (a) | 225.0 | (a) | 1,816.4 |
| Total, Middle Atlantic | 1,816.7 | 24.5 | (a) | (a) | 256.6 | (a) | 2,097.9 |
| Softwoods | 273.5 | 20.3 | 43.8 | 2.1 | 105.6 | (a) | 445.1 |
| Hardwoods | 918.6 | 71.0 | 55.5 | 29.0 | 184.1 | 3.7 | 1,261.9 |
| Total, Lake States | 1,192.0 | 91.3 | 99.3 | 31.1 | 289.7 | 3.7 | 1,707.0 |
| Softwoods | 49.6 | 2.7 | 3.2 | 0.1 | 18.3 | (a) | 74.0 |
| Hardwoods | 1,062.2 | 78.5 | 30.3 | 10.8 | 266.2 | 2.1 | 1,450.1 |
| Total, Central States | 1,111.8 | 81.2 | 33.4 | 10.9 | 284.5 | 2.1 | 1,524.0 |
| Softwoods | 1,090.9 | 34.0 | 46.9 | 2.2 | 234.0 | (a) | 1,408.1 |
| Hardwoods | 4,228.8 | 183.2 | 85.8 | 39.8 | 777.9 | 5.8 | 5,321.2 |
| Total, North | 5,319.6 | 217.2 | 132.7 | 42.0 | 1,011.9 | 5.8 | 6,729.3 |

(continued)

Table 7.--Total forest biomass on timberland in the United States, by major species group, section, region, and class of material, 1987, continued

| Species group, section, and region | Biomass of live sound and cull trees | Other live tree biomass | | Other forest biomass | | | Total biomass |
|--|--|-------------------------|---------|----------------------|----------|--------|------------------|
| | | Stumps | Foliage | Seedlings | Saplings | Shrubs | |
| ----- (Million dry tons) ----- | | | | | | | |
| Softwoods | 572.0 | 38.9 | 35.0 | 0.4 | 56.3 | (a) | 702.6 |
| Hardwoods | 1,210.5 | 94.7 | 68.7 | 0.6 | 241.8 | (a) | 1,616.3 |
| Total, South Atlantic | 1,782.5 | 133.6 | 103.7 | 1.0 | 298.1 | (a) | 2,318.9 |
| Softwoods | 552.8 | 41.3 | 32.1 | 0.5 | 65.0 | (a) | 691.7 |
| Hardwoods | 575.2 | 46.1 | 34.1 | 0.4 | 133.0 | (a) | 788.8 |
| Total, East Gulf | 1,128.0 | 87.4 | 66.2 | 0.9 | 198.0 | (a) | 1,480.5 |
| Softwoods | 546.3 | 27.2 | 18.7 | (a) | 55.2 | (a) | 647.4 |
| Hardwoods | 1,372.5 | 51.8 | 26.0 | (a) | 252.6 | (a) | 1,702.9 |
| Total, Central Gulf | 1,918.8 | 79.0 | 44.7 | (a) | 307.8 | (a) | 2,350.3 |
| Softwoods | 600.8 | 22.9 | 22.5 | (a) | 51.0 | (a) | 697.2 |
| Hardwoods | 1,115.6 | 53.0 | 34.4 | (a) | 190.7 | (a) | 1,393.7 |
| Total, West Gulf | 1,716.4 | 75.9 | 56.9 | (a) | 241.7 | (a) | 2,090.9 |
| Softwoods | 2,271.9 | 130.3 | 108.3 | 0.9 | 227.5 | (a) | 2,738.9 |
| Hardwoods | 4,273.8 | 245.6 | 163.2 | 1.0 | 818.1 | (a) | 5,501.7 |
| Total, South | 6,545.7 | 375.9 | 271.5 | 1.9 | 1,045.6 | (a) | 8,240.6 |

(continued)

Table 7.--Total forest biomass on timberland in the United States, by major species group, section, region, and class of material, 1987, continued

| Species group, section, and region | Biomass of live sound and cull trees | Other live tree biomass | | | Other forest biomass | | | Total biomass |
|--|--|-------------------------|---------|-----------|----------------------|--------|--|------------------|
| | | Stumps | Foliage | Seedlings | Saplings | Shrubs | | |
| | | | | | | | | |
| ----- (Million dry tons) ----- | | | | | | | | |
| Softwoods | 1,285.6 | (a) | (a) | (a) | 80.9 | (a) | | 1,366.5 |
| Hardwoods | 19.8 | (a) | (a) | (a) | 1.4 | (a) | | 21.2 |
| Total, N. Rocky Mountain | 1,305.5 | (a) | (a) | (a) | 82.3 | (a) | | 1,387.8 |
| Softwoods | 559.0 | (a) | (a) | (a) | 24.6 | (a) | | 583.6 |
| Hardwoods | 77.7 | (a) | (a) | (a) | 4.5 | (a) | | 82.2 |
| Total, S. Rocky Mountain | 636.7 | (a) | (a) | (a) | 29.1 | (a) | | 665.8 |
| Softwoods | 2,789.3 | (a) | (a) | (a) | 128.6 | (a) | | 2,917.8 |
| Hardwoods | 458.0 | (a) | (a) | (a) | 56.6 | (a) | | 514.5 |
| Total, Pacific Coast | 3,247.2 | (a) | (a) | (a) | 185.1 | (a) | | 3,432.3 |
| Softwoods | 659.8 | 16.0 | 8.0 | 1.4 | 35.3 | (a) | | 720.4 |
| Hardwoods | 119.3 | 2.1 | 1.0 | 0.3 | 51.9 | 75.9 | | 250.6 |
| Total, Alaska and Hawaii | 779.1 | 18.0 | 9.0 | 1.7 | 87.3 | 75.9 | | 971.0 |
| Softwoods | 5,293.8 | 16.0 | 8.0 | 1.4 | 269.4 | (a) | | 5,588.4 |
| Hardwoods | 674.8 | 2.1 | 1.0 | 0.3 | 114.4 | 75.9 | | 868.5 |
| Total, West | 5,968.6 | 18.0 | 9.0 | 1.7 | 383.8 | 75.9 | | 6,457.0 |
| Softwoods | 8,656.5 | 180.3 | 163.2 | 4.5 | 730.9 | (a) | | 9,735.4 |
| Hardwoods | 9,177.4 | 430.9 | 250.0 | 41.2 | 1,710.4 | 81.6 | | 11,691.5 |
| Total, all regions | 17,833.9 | 611.2 | 413.2 | 45.6 | 2,441.3 | 81.6 | | 21,426.9 |

^a Data unavailable.

Table 8.--Tree biomass per acre on forest land in the United States, by major species group, section, region, and land class, 1987

| Species group, section, and region | Timberland | | | | | Other forest land | Total forest land |
|--|---------------------------------|-----------|-------|-----------------------|-------|----------------------|----------------------|
| | Live sound trees | | | Live cull trees | Total | | |
| | Tree boles | Tree tops | Total | | | | |
| | ----- (Dry tons per acre) ----- | | | | | | |
| Softwoods | 10.90 | 4.09 | 14.99 | 1.52 | 16.51 | (a) | (a) |
| Hardwoods | 13.10 | 4.91 | 18.01 | 3.49 | 21.49 | (a) | (a) |
| Total, New England | 24.00 | 9.00 | 32.99 | 5.00 | 38.00 | (a) | (a) |
| Softwoods | 3.59 | 1.20 | 4.79 | 0.30 | 5.09 | (a) | (a) |
| Hardwoods | 22.75 | 7.31 | 30.05 | 2.28 | 32.34 | (a) | (a) |
| Total, Middle Atlantic | 26.33 | 8.51 | 34.84 | 2.58 | 37.42 | (a) | (a) |
| Softwoods | 4.58 | 0.92 | 5.51 | 0.26 | 5.76 | (a) | (a) |
| Hardwoods | 13.18 | 4.09 | 17.28 | 2.09 | 19.36 | (a) | (a) |
| Total, Lake States | 17.77 | 5.02 | 22.78 | 2.34 | 25.13 | (a) | (a) |
| Softwoods | 0.82 | 0.27 | 1.08 | 0.08 | 1.16 | (a) | (a) |
| Hardwoods | 15.35 | 4.39 | 19.74 | 5.16 | 24.90 | (a) | (a) |
| Total, Central States | 16.17 | 4.66 | 20.82 | 5.24 | 26.06 | (a) | (a) |
| Softwoods | 4.53 | 1.42 | 5.95 | 0.46 | 6.41 | (a) | (a) |
| Hardwoods | 16.44 | 5.24 | 21.67 | 3.17 | 24.85 | (a) | (a) |
| Total, North | 20.96 | 6.66 | 27.62 | 3.63 | 31.25 | (a) | (a) |

(continued)

Table 8.--Tree biomass per acre on forest land in the United States, by major species group, section, region, and land class, 1987,
continued

| Species group, section, and region | Timberland | | | | | Other forest land | Total forest land |
|--|---------------------------------|-----------|-------|-----------------------|-------|----------------------|----------------------|
| | Live sound trees | | | Live cull trees | Total | | |
| | Tree boles | Tree tops | Total | | | | |
| | ----- (Dry tons per acre) ----- | | | | | | |
| Softwoods | 11.06 | 1.24 | 12.30 | 0.14 | 12.44 | 0.60 | 12.15 |
| Hardwoods | 20.19 | 3.01 | 23.21 | 3.12 | 26.33 | 7.00 | 25.86 |
| Total, South Atlantic | 31.25 | 4.25 | 35.51 | 3.26 | 38.77 | 7.60 | 38.02 |
| Softwoods | 12.69 | 1.50 | 14.19 | 0.12 | 14.31 | 1.50 | 13.68 |
| Hardwoods | 10.86 | 1.64 | 12.50 | 2.40 | 14.89 | 1.80 | 14.25 |
| Total, East Gulf | 23.55 | 3.13 | 26.69 | 2.52 | 29.21 | 3.30 | 27.93 |
| Softwoods | 8.42 | 1.59 | 10.01 | 0.59 | 10.60 | 0.00 | 10.60 |
| Hardwoods | 14.95 | 5.06 | 20.01 | 6.59 | 26.60 | 0.00 | 26.60 |
| Total, Central Gulf | 23.39 | 6.66 | 30.05 | 7.20 | 37.24 | 0.00 | 37.24 |
| Softwoods | 10.41 | 1.84 | 12.25 | 0.45 | 12.70 | 1.60 | 12.60 |
| Hardwoods | 12.46 | 4.54 | 17.01 | 6.51 | 23.50 | 19.00 | 21.60 |
| Total, West Gulf | 22.86 | 6.37 | 29.24 | 6.95 | 36.19 | 20.60 | 35.01 |
| Softwoods | 10.49 | 1.55 | 12.03 | 0.34 | 12.38 | 1.41 | 11.97 |
| Hardwoods | 14.77 | 3.69 | 18.46 | 4.82 | 23.28 | 12.13 | 22.87 |
| Total, South | 25.26 | 5.24 | 30.50 | 5.16 | 35.66 | 13.54 | 34.85 |

(continued)

Table 8.--Tree biomass per acre on forest land in the United States, by major species group, section, region, and land class, 1987, continued

| Species group, section, and region | Timberland | | | | | Other forest land | Total forest land |
|--|------------------|-----------|-------|-----------------------|-------|----------------------|----------------------|
| | Live sound trees | | | Live cull trees | Total | | |
| | Tree boles | Tree tops | Total | | | | |
| ----- (Dry tons per acre) ----- | | | | | | | |
| Softwoods | 24.60 | 7.25 | 31.84 | 6.42 | 38.26 | (a) | (a) |
| Hardwoods | 0.39 | 0.10 | 0.48 | 0.11 | 0.59 | (a) | (a) |
| Total, N. Rocky Mountain | 24.98 | 7.34 | 32.32 | 6.53 | 38.85 | (a) | (a) |
| Softwoods | 15.33 | 5.39 | 20.72 | 2.57 | 23.28 | (a) | (a) |
| Hardwoods | 2.13 | 0.45 | 2.58 | 0.66 | 3.24 | (a) | (a) |
| Total, S. Rocky Mountain | 17.46 | 5.84 | 23.29 | 3.23 | 26.52 | (a) | (a) |
| Softwoods | 42.28 | 7.24 | 49.52 | 0.91 | 50.43 | (a) | (a) |
| Hardwoods | 6.19 | 1.29 | 7.48 | 0.80 | 8.28 | (a) | (a) |
| Total, Pacific Coast | 48.47 | 8.53 | 57.00 | 1.71 | 58.71 | (a) | (a) |
| Softwoods | 31.80 | 6.78 | 38.58 | 1.50 | 40.08 | (a) | (a) |
| Hardwoods | 5.48 | 1.19 | 6.67 | 0.58 | 7.25 | (a) | (a) |
| Total, Alaska and Hawaii | 37.28 | 7.96 | 45.25 | 2.08 | 47.33 | (a) | (a) |
| Softwoods | 31.35 | 6.84 | 38.19 | 2.73 | 40.92 | (a) | (a) |
| Hardwoods | 3.84 | 0.81 | 4.65 | 0.57 | 5.22 | (a) | (a) |
| Total, West | 35.19 | 7.65 | 42.84 | 3.29 | 46.13 | (a) | (a) |
| Softwoods | 13.98 | 2.92 | 16.90 | 1.02 | 17.92 | (a) | (a) |
| Hardwoods | 12.43 | 3.47 | 15.90 | 3.10 | 19.00 | (a) | (a) |
| Total, all regions | 26.41 | 6.39 | 32.79 | 4.12 | 36.91 | (a) | (a) |

^a Data unavailable.

